arXiv:1607.02591v1 [math.RA] 9 Jul 2016

On the decomposition of metabolic involutions

A.-H. Nokhodkar

Abstract

The problem of whether a metabolic idempotent of a central simple algebra with involution is contained in an invariant quaternion subalgebra is investigated. As an application, the similar problem is studied for skew-symmetric elements whose squares lie in the square of the underlying field.

Mathematics Subject Classification: 16W10, 16K20, 16K50.
Keywords: Central simple algebra, Division algebra, Involution.

1 Introduction

A hyperbolic involution on a central simple algebra is an involution which is adjoint to a hyperbolic hermitian form. It is known that hyperbolic involutions are decomposable (see [2, (2.2)] and [7, Ch. II, Exercise 2]). Indeed, using the ideas of [2], one can show that every hyperbolic idempotent of a central simple algebra with involution (A, σ) lies in a σ-invariant quaternion subalgebra (see (4.1) below). A weaker condition for involutions is *metabolicity*. Metabolic involutions were introduced in [3] and studied in more detail in [5]. According to [5, (4.8)], metabolic involutions are adjoint to metabolic hermitian forms.

A decomposition problem concerning metabolic involutions is that whether a metabolic idempotent of a central simple algebra with involution (A, σ) is contained in a σ-invariant quaternion subalgebra. Since every metabolic idempotent e satisfies $(e - \sigma(e))^2 = 1$, such a quaternion subalgebra would also contain a skew-symmetric element whose square equals 1. Hence, a relevant decomposition problem arises as follows: given a skew-symmetric element $u \in A$ satisfying $u^2 = 1$ (or more generally, $u^2 \in F^2$), is there any σ-invariant quaternion subalgebra of A containing u?

The aim of this work is to study the aforementioned decomposition problems in arbitrary characteristic. We start with some general observations on idempotents and square-central elements in a central simple algebra. Let A be a central simple algebra over a field F and let $u \in A$ be an element satisfying $u^2 = \lambda^2$ for some $\lambda \in F^\times$. In [1, (4.1)], it was shown that if $\operatorname{char} F \neq 2$, then u is contained in a quaternion subalgebra of A if and only if $\dim_F(u + \lambda)A = \dim_F(u - \lambda)A$. Using similar methods we shall generalize this result in (3.2) to arbitrary characteristic, including $\lambda = 0$.

In §4 we study metabolic idempotents of a central simple algebra (A, σ). The main result is (4.4), which asserts that a metabolic idempotent $e \in A$ lies in a σ-invariant quaternion subalgebra if and only if it is hyperbolic or $\dim_F e\sigma(e)A = \frac{1}{2}\dim_F A$. Using this, we will see in (4.5) that in contrast with the case of hyperbolic idempotents, there exists a metabolic idempotent with

respect to an involution σ (over a field of arbitrary characteristic), which is not contained in any σ-invariant quaternion subalgebra.

In the last section we use the decomposition criterion (4.4) to study skew-symmetric elements of A whose squares lie in F^2. Consider an element $u \in A$ satisfying $u^2 = \lambda^2$ for some $\lambda \in F$ and $\dim_F (u+\lambda)A = \frac{1}{2}\dim_F A$. As we shall see in (5.1), if $u \in \mathrm{Alt}(A,\sigma)$, $\sigma(\lambda) = \lambda$ and $\lambda \neq 0$, then there exists a metabolic idempotent $e \in A$ such that $u = e - \sigma(e)$. An important application of this result is (5.3) which asserts that if $\sigma(u) = -u$, then there exists a σ-invariant quaternion subalgebra of A containing u, except for the case where $\mathrm{char}\, F = 2$ and σ is orthogonal. Finally, this exceptional case is treated in (5.5) and (5.6).

2 Preliminaries

Let A be a central simple algebra over a field F and let σ be an *involution* on A, i.e., an anti-automorphism satisfying $\sigma^2 = \mathrm{id}$. We say that σ is *of the first kind* if $\sigma|_F = \mathrm{id}$. Otherwise, σ is said to be *of the second kind*. Involutions of the second kind are also called *unitary* involutions. After scalar extension to a splitting field of A, every involution σ of the first kind becomes adjoint to a symmetric or skew-symmetric bilinear form. If this bilinear form is alternating, we say that σ is *symplectic*. Otherwise, σ is called *orthogonal*. The set of all *symmetric* elements of an algebra with involution (A,σ), i.e., elements fixed by σ, is denoted by $\mathrm{Sym}(A,\sigma)$. We also use the notations

$$\mathrm{Symd}(A,\sigma) = \{x + \sigma(x) \mid x \in A\} \quad \text{and} \quad \mathrm{Alt}(A,\sigma) = \{x - \sigma(x) \mid x \in A\}.$$

An algebra with involution (A,σ) (or simply the involution σ) is called *hyperbolic* if it has a *hyperbolic idempotent* with respect to σ, i.e., an idempotent $e \in A$ satisfying $\sigma(e) = 1 - e$. Note that for every hyperbolic idempotent e we have $\dim_F eA = \frac{1}{2}\dim_F A$ (see [7, p. 75]). An idempotent $e \in A$ is called *metabolic* if (1) $\sigma(e)e = 0$ and (2) $\dim_F eA = \frac{1}{2}\dim_F A$. Either of these two conditions can be equivalently substituted with $(1-e)(1-\sigma(e)) = 0$, as observed in [3] and [5]. The pair (A,σ) (or simply the involution σ) is called *metabolic* if there exists a metabolic idempotent $e \in A$ with respect to σ. Every hyperbolic idempotent is metabolic, but there exist metabolic idempotents (in all characteristics) that are not hyperbolic. However, if e is a metabolic idempotent and there exists an element $x \in A$ with $x + \sigma(x) = 1$, then the element $e' := e - ex\sigma(e)$ is a hyperbolic idempotent of (A,σ) (see [3, (A.3)]). Combining this observation with (2.1) below, one concludes that an involution σ is hyperbolic if and only if it is metabolic, except for the case where $\mathrm{char}\, F = 2$ and σ is orthogonal. Note that in view of [7, (2.6 (2))], hyperbolic involutions can never exist in this exceptional case.

Lemma 2.1. *Let (A,σ) be a central simple algebra with involution over a field F. Then there exists $x \in A$ such that $x + \sigma(x) = 1$, except for the case where $\mathrm{char}\, F = 2$ and σ is orthogonal.*

Proof. If $\mathrm{char}\, F \neq 2$, take $x = \frac{1}{2}$. Otherwise, σ is either symplectic or unitary and the existence of such an element follows from [7, (2.6 (2)) and (2.17)]. □

3 Idempotents and square-central elements

Let D be a central division algebra over a field F. We write $M_n(D)$ for the algebra of all $n \times n$ matrices with entries in D. The identity matrix in $M_n(D)$ is denoted by I_n. Also, for $\alpha \in D$, we denote by $J_n(\alpha)$ the $n \times n$ *Jordan block* whose diagonal and superdiagonal entries equal to α and 1 respectively and all other entries are zero.

Our first result is easily deduced from [4, Prop. 6 and Thm. 7].

Lemma 3.1. *Let V be a finite-dimensional right vector space over D and let $x \in \mathrm{End}_D(V)$.*

(1) If $x^2 = \lambda^2$ for some $\lambda \in F$, then there exist integers m, n, k and a basis of V with respect to which the matrix of x is a direct sum of λI_m, $-\lambda I_n$ and k Jordan blocks $J_2(\lambda)$ (the blocks $J_2(\lambda)$ appear only if $\mathrm{char}\, F = 2$ or $\lambda = 0$).

(2) If $x^2 = x$, then there exist an integer m and a basis of V with respect to which the matrix of x is a direct sum of I_m and a zero matrix.

The next result generalizes [1, (4.1)] to arbitrary characteristic. Recall that a *quaternion algebra* over a field F is a central simple F-algebra of degree 2.

Proposition 3.2. ([1, (4.1)]) *Let A be a central simple algebra over a field F and let $u \in A \setminus F$ such that $u^2 = \lambda^2 \in F^2$ for some $\lambda \in F$. There exists a split quaternion F-subalgebra of A containing u if and only if $\dim_F(\lambda + u)A = \frac{1}{2}\dim_F A$.*

Proof. If $\mathrm{char}\, F \neq 2$ and $\lambda \neq 0$, the result is easily deduced from [1, (4.1)], hence let $\mathrm{char}\, F = 2$ or $\lambda = 0$. Suppose that u is contained in a split quaternion subalgebra of A. We identify this subalgebra with $\mathrm{End}_F(W)$ for some 2-dimensional vector space W over F. As $u \notin F$, by (3.1 (1)) there exists a basis of W with respect to which the matrix of u equals the Jordan block $J_2(\lambda)$. Since $\lambda = 0$ or $\mathrm{char}\, F = 2$, $\lambda + u$ is represented in this basis by $J_2(0)$. Thus, $\dim_F(\lambda + u)\,\mathrm{End}_F(W) = 2 = \frac{1}{2}\dim_F \mathrm{End}_F(W)$, which leads to the desired equality.

To prove the converse, observe first that the conditions $u \notin F$ and $u^2 \in F^2$ imply that A is not a division algebra. Choose a division F-algebra D, Brauer equivalent to A. Then we may identify $A = \mathrm{End}_D(V)$, where V is a right D-vector space. Since $u^2 = \lambda^2$, by (3.1 (1)) there exist an integer r and a basis $\mathcal{B}$ of V with respect to which the matrix of u is a direct sum of λI_r and the Jordan blocks $J_2(\lambda)$. Let s be the number of blocks $J_2(\lambda)$ in this decomposition, so that $r + 2s = \dim_D V$, where $\dim_D V$ denotes the right dimension of V over D. Since $\lambda = 0$ or $\mathrm{char}\, F = 2$, the matrix of $\lambda + u$ in $\mathcal{B}$ is a direct sum of s blocks $J_2(0)$ and an $r \times r$ zero matrix, hence $\dim_D(\lambda + u)A = s(r + 2s)$. The equality $\dim_D(\lambda + u)A = \frac{1}{2}\dim_D A = \frac{1}{2}(r + 2s)^2$ then implies that $r = 0$. Thus, u is a direct sum of the Jordan blocks $J_2(\lambda)$ (in particular, $\dim_D V$ is even). Let $v \in \mathrm{End}_D(V)$ be the endomorphism which is represented in $\mathcal{B}$ by direct sums of matrices of the form

$$\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Then the elements 1, u, v and uv span a subalgebra of $A = \mathrm{End}_D(V)$, isomorphic to $M_2(F)$. □

Lemma 3.3. *Let A be a central simple algebra over a field F and let $e \in A$ be an idempotent. Suppose that there exists a nonzero element $u \in A$ satisfying $u^2 = \lambda^2$ for some $\lambda \in F$, $ue = \lambda e$ and $e(\lambda + u) = \lambda + u$. If $\dim_F eA = \dim_F e(u - \lambda)A = \frac{1}{2}\dim_F A$, then there exists an element $w \in A$ such that $w^2 = 0$, $ew = 0$, $we = w$, $uw = e - \lambda w$ and $wu = \lambda w - e + 1$. Furthermore, the elements 1, e, u and w span a split quaternion subalgebra of A.*

Proof. We may identify $A = \mathrm{End}_D(V)$, where D is a division F-algebra Brauer equivalent to A and V is a right vector space over D. Let $n = \dim_D V$. The condition $\dim_F eA = \frac{1}{2}\dim_F A$ implies that e is a nontrivial idempotent. Hence A is not a division algebra, i.e., $n > 1$. By (3.1) there exist an integer m and a D-basis $\mathcal{B}$ of V, with respect to which e is represented by the matrix

$$e = \begin{pmatrix} I_m & 0 \\ 0 & 0 \end{pmatrix} \in M_n(D).$$

Since $\dim_F eA = \frac{1}{2}\dim_F A$, we have $m = \frac{n}{2}$. Let

$$\begin{pmatrix} U & W \\ X & Y \end{pmatrix},$$

denote the matrix of u in $\mathcal{B}$, where $U, W, X, Y \in M_m(D)$. The relations $ue = \lambda e$ and $e(\lambda + u) = \lambda + u$ imply that $U = \lambda I_m$, $X = 0$ and $Y = -\lambda I_m$, i.e.,

$$u = \begin{pmatrix} \lambda I_m & W \\ 0 & -\lambda I_m \end{pmatrix},$$

which yields

$$e(u - \lambda) = \begin{pmatrix} 0 & W \\ 0 & 0 \end{pmatrix}.$$

It follows that

$$e(u - \lambda)M_n(D) = \left\{ \begin{pmatrix} WM & WN \\ 0 & 0 \end{pmatrix} \mid M, N \in M_m(D) \right\}.$$

The equality $\dim_F e(u - \lambda)A = \frac{1}{2}\dim_F A$ implies that W has a right inverse, hence it is invertible by [6, §19, Thm. 3]. A direct calculation then shows that the element

$$w = \begin{pmatrix} 0 & 0 \\ W^{-1} & 0 \end{pmatrix},$$

has the desired properties. □

4 Metabolic and hyperbolic idempotents

Let $(D, {}^-)$ be a division algebra with involution over a field F and let V be a right D-vector space. Denote by $V^* = \mathrm{Hom}_D(V, D)$, the dual space of V, with a right D-vector space structure $v^* d(v) = \bar{d}v^*(v)$ for $v \in V$, $v^* \in V^*$ and $d \in D$. For every $\lambda \in F$ with $\lambda\bar{\lambda} = 1$, define the λ-hermitian form h_λ on $H_\lambda(V) := V^* \oplus V$ via

$$h_\lambda((v_1^*, v_1), (v_2^*, v_2)) = v_1^*(v_2) + \lambda\overline{v_2^*(v_1)}.$$

A λ-hermitian space is called *hyperbolic* if it is isometric to $H_\lambda(V)$ for some right vector space V over D.

The following result is implicitly contained in [2, pp. 466-468] for characteristic different from 2.

Theorem 4.1. ([2]) *Let (A, σ) be a central simple algebra with involution over a field F. For every hyperbolic idempotent $e \in A$ there exists a σ-invariant split quaternion subalgebra of A containing e.*

Proof. Let D be a division algebra Brauer-equivalent to A and let $^-\colon D \to D$ be an involution on D such that $^-|_F = \sigma|_F$. We may identify

$$(A, \sigma) = (\mathrm{End}_V(D), \sigma_h),$$

where V is a right vector space over D, λ is an element in D with $\lambda\bar{\lambda} = 1$, $h : V \times V \to D$ is a non-degenerate λ-hermitian form and σ_h is the adjoint involution of $\mathrm{End}_D(V)$ with respect to h. Let $S = \mathrm{im}\, e \subseteq V$ and $T = \mathrm{im}\, \sigma(e) \subseteq V$ be the images of e and $\sigma(e)$. Define a map $\varphi : S \to T^*$ via $\varphi(s)(t) = h(s, t)$ for $s \in S$ and $t \in T$. As observed in the proof of [2, (2.1)], the map $\varphi \oplus \mathrm{id}$ defines an isometry between $(V, h) = (S \oplus T, h)$ and $(H_\lambda(T), h_\lambda)$. Considering this isomorphism as an identification, we have

$$e(t^*, t) = (t^*, 0) \quad \text{and} \quad \sigma(e)(t^*, t) = (0, t) \quad \text{for } t^* \in T^* \text{ and } t \in T.$$

Let $\ell : T \times T \to D$ be a non-degenerate 1-hermitian form and let $\theta : T \to T^*$ be the isomorphism of right D-vector spaces defined by $\theta(t_1)(t_2) = \ell(t_1, t_2)$ for $t_1, t_2 \in T$. Define $u, v \in A = \mathrm{End}_D(V)$ via

$$u(t^*, t) = (\theta(t), 0) \quad \text{and} \quad v(t^*, t) = (0, \theta^{-1}(t^*)).$$

As observed in the proof of [2, (2.2)], the elements e, $\sigma(e)$, u and v span a σ-invariant subalgebra of A, isomorphic to $M_2(F)$. □

Lemma 4.2. *Let (A, σ) be a central simple algebra with involution over a field F and let $e \in A$ be a hyperbolic idempotent. Suppose that there exists $u \in A$ satisfying $u^2 = 0$, $\dim_F uA = \frac{1}{2}\dim_F A$, $ue = 0$ and $eu = u$. If $\sigma(u) = \pm u$, then there exists a σ-invariant quaternion subalgebra of A containing u and e.*

Proof. Since e is a hyperbolic idempotent, we have $\dim_F eA = \frac{1}{2}\dim_F A$. Hence, using (3.3) with $\lambda = 0$ one can find $w \in A$ such that

$$w^2 = 0, \quad ew = 0, \quad we = w, \quad uw = e \quad \text{and} \quad wu = 1 - e = \sigma(e).$$

Also, the F-algebra $Q \subseteq A$ spanned by 1, e, u and w is a quaternion algebra. We claim that $\sigma(Q) = Q$. Since $\sigma(u) = \pm u \in Q$ and $\sigma(e) = 1 - e \in Q$, it suffices to show that $\sigma(w) \in Q$. Keeping the notations of the proof of (3.3) with $\lambda = 0$ we have

$$u = \begin{pmatrix} 0 & W \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \sigma(e) = 1 - e = \begin{pmatrix} 0 & 0 \\ 0 & I_m \end{pmatrix},$$

where $W \in M_m(D)$ is invertible. Write

$$\sigma(w) = \begin{pmatrix} X & Y \\ Z & T \end{pmatrix},$$

for some $X, Y, Z, T \in M_m(D)$. Suppose first that $\sigma(u) = u$. Applying σ on $uw = e$, we get $\sigma(w)u = \sigma(e)$, which implies that $X = 0$ and $Z = W^{-1}$. Similarly, applying σ on the equality $wu = \sigma(e)$, we obtain $u\sigma(w) = e$, hence $T = 0$. Finally, as $w^2 = 0$, we have $\sigma(w)^2 = 0$, which leads to $Y = 0$. It follows that

$$\sigma(w) = \begin{pmatrix} 0 & 0 \\ W^{-1} & 0 \end{pmatrix} = w \in Q.$$

If $\sigma(u) = -u$, a similar argument shows that $\sigma(w) = -w \in Q$. □

Lemma 4.3. *Let (A, σ) be a central simple algebra with involution over a field F and let $e \in A$ be a metabolic idempotent.*

(1) e is a hyperbolic idempotent if and only if $e\sigma(e) = 0$.

(2) For every $x \in A$, the element $e' := e - ex\sigma(e)$ is a metabolic idempotent.

Proof. The first statement follows from the equality $1 - e - \sigma(e) + e\sigma(e) = 0$. To prove (2) note that the equality $\sigma(e)e = 0$ implies that e' is an idempotent and $\sigma(e')e' = 0$. Also, since $ee' = e'$ and $e'e = e$, we have $eA = e'A$. It follows that $\dim_F e'A = \dim_F eA = \frac{1}{2}\dim_F A$. Hence e' is a metabolic idempotent. □

Theorem 4.4. *Let (A, σ) be a central simple algebra with involution over a field F and let $e \in A$ be a metabolic idempotent. There exists a σ-invariant split quaternion subalgebra of A containing e if and only if either e is hyperbolic or $\dim_F e\sigma(e)A = \frac{1}{2}\dim_F A$.*

Proof. Set $u = e\sigma(e)$. Suppose that there exists a σ-invariant quaternion subalgebra of A containing e. If e is not hyperbolic, then we have $u \neq 0$ by (4.3 (1)). As $u^2 = 0$, using (3.2) with $\lambda = 0$ we obtain $\dim_F uA = \frac{1}{2}\dim_F A$. Conversely, if e is hyperbolic the result follows from (4.1). Suppose that $\dim_F uA = \frac{1}{2}\dim_F A$. We proceed similar to the proof of (4.2). We have $u^2 = 0$, $ue = 0$, $eu = u$ and $\dim_F euA = \dim_F uA = \frac{1}{2}\dim_F A$. Also, $\dim_F eA = \frac{1}{2}\dim_F A$, since e is a metabolic idempotent. Thus, one can use (3.3) with $\lambda = 0$ to find $w \in A$ such that

$$w^2 = 0, \quad ew = 0, \quad we = w, \quad uw = e \quad \text{and} \quad wu = 1 - e.$$

By (3.3), the F-algebra $Q \subseteq A$ spanned by 1, e, u and w is a quaternion algebra. We claim that $\sigma(Q) = Q$. We have $\sigma(u) = u \in Q$. Also, the equality $(1 - e)(1 - \sigma(e)) = 0$ implies that $\sigma(e) = 1 - e + u \in Q$. Thus, it remains to show that $\sigma(w) \in Q$. Keep the notation of the proof of (3.3) with $\lambda = 0$, so that

$$u = \begin{pmatrix} 0 & W \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \sigma(e) = 1 - e + u = \begin{pmatrix} 0 & W \\ 0 & I_m \end{pmatrix},$$

where W is an invertible matrix in $M_m(D)$. Write

$$\sigma(w) = \begin{pmatrix} X & Y \\ Z & T \end{pmatrix},$$

for some $X, Y, Z, T \in M_m(D)$. Applying σ on $uw = e$, we get $\sigma(w)u = \sigma(e)$, which leads to $X = I_m$ and $Z = W^{-1}$. Also, applying σ on $wu = 1 - e$, we obtain $u\sigma(w) = 1 - \sigma(e)$, so $T = -I_m$. Finally, the equality $w^2 = 0$ implies that $\sigma(w)^2 = 0$, hence $Y = -W$. It follows that

$$\sigma(w) = \begin{pmatrix} I_m & -W \\ W^{-1} & -I_m \end{pmatrix} = w + 2e - u - 1 \in Q.$$

□

The next example shows the existence of a metabolic idempotent in a central simple algebra with involution over a field of arbitrary characteristic which is not contained in any invariant quaternion subalgebra.

Example 4.5. Let F be a field and let t be the transpose involution on $M_4(F)$. Set $\sigma = \mathrm{Int}(u) \circ t$, where

$$u = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix},$$

and $\mathrm{Int}(u)$ is the inner automorphism of $M_4(F)$ induced by u, i.e., $\mathrm{Int}(u)(x) = uxu^{-1}$ for $x \in M_4(F)$. Since u is a symmetric matrix, by [7, (2.7 (1))], σ is an involution of the first kind on $M_4(F)$. A straightforward calculation shows that the element

$$e := \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 \end{pmatrix},$$

is a metabolic idempotent with respect to σ. We also have

$$e\sigma(e) = \begin{pmatrix} 1 & -1 & 1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & 1 & -1 \end{pmatrix},$$

hence e is not hyperbolic and $\dim_F e\sigma(e)M_4(F) = 4 \neq \frac{1}{2}\dim_F M_4(F)$. By (4.4) there is no σ-invariant quaternion subalgebra of $M_4(F)$ containing e.

5 Applications to square-central elements

Throughout this section, (A, σ) denotes a central simple algebra with involution over a field F and $u \in A \setminus F$ is an element satisfying $u^2 = \lambda^2$ for some $\lambda \in F$ with $\sigma(\lambda) = \lambda$ and

$$\dim_F(\lambda + u)A = \frac{1}{2}\dim_F A. \tag{1}$$

Our purpose is to find some sufficient conditions on u to be contained in a σ-invariant quaternion subalgebra of A. Note that in view of (3.2), the condition (1) is necessary for this aim. Also, if σ is of the first kind, then the equality $\sigma(\lambda) = \lambda$ holds automatically.

For a right ideal I of A let $I^\perp = \{x \in A \mid \sigma(x)y = 0 \text{ for } y \in I\}$. Then $I^\perp$ is also a right ideal of A. Moreover, according to [7, (6.2)] we have

$$\dim_F I + \dim_F I^\perp = \dim_F A.$$

Proposition 5.1. *Suppose that $\sigma(u) = -u$. Then*

(1) σ is metabolic.

(2) There exists a metabolic idempotent $e \in A$ such that $e(\lambda + u) = \lambda + u$ and $ue = \lambda e$.

(3) If $\lambda \neq 0$ and $u \in \mathrm{Alt}(A,\sigma)$, then there exists a metabolic idempotent $e' \in A$ such that $\lambda^{-1}u = e' - \sigma(e')$. In addition, if $\operatorname{char} F = 2$, such an idempotent satisfies $\dim_F e'\sigma(e') = \frac{1}{2}\dim_F A$. If $\operatorname{char} F \neq 2$, the idempotent $e' \in A$ can be chosen to be hyperbolic.

Proof. Consider the right ideal $I := (\lambda + u)A$ of A. Since $\sigma(\lambda + u) = \lambda - u$, we have $\sigma(I)I = 0$. By dimension count we obtain $I = I^\perp$, hence σ is metabolic by [5, (4.4)]. This proves (1).

(2) By [7, (1.13)] there exists an idempotent $e \in A$ such that $I = eA$. As $I = I^\perp$, we have $\sigma(e)e = 0$. We also have $\dim_F eA = \dim_F(\lambda+u)A = \frac{1}{2}\dim_F A$, hence e is a metabolic idempotent. Since $(\lambda + u)A = eA$, there exist $x, y \in A$ such that $\lambda + u = ex$ and $e = (\lambda + u)y$. Multiplying the first equality on the left by e, we get $e(\lambda+u) = e^2x = ex = \lambda+u$. Similarly, multiplying the second equality on the left by u, we obtain $ue = \lambda e$.

(3) Applying σ on $e(\lambda + u) = \lambda + u$ we get $(\lambda - u)\sigma(e) = \lambda - u$, i.e., $u\sigma(e) = \lambda\sigma(e) + u - \lambda$. Also, $eu = e(\lambda + u) - \lambda e = \lambda + u - \lambda e$, hence

$$eu\sigma(e) = e(\lambda\sigma(e) + u - \lambda) = \lambda e\sigma(e) + eu - \lambda e = \lambda e\sigma(e) + \lambda + u - 2\lambda e. \quad (2)$$

Now, write $\lambda^{-1}u = x - \sigma(x)$ for some $x \in A$ and set $e' = e - e\sigma(x)\sigma(e)$. By (4.3 (2)), e' is a metabolic idempotent. Using the relation (2) we get

$$\begin{aligned} e' - \sigma(e') &= e - e\sigma(x)\sigma(e) - \sigma(e) + ex\sigma(e) = e - \sigma(e) + e(x - \sigma(x))\sigma(e) \\ &= e - \sigma(e) + \lambda^{-1}eu\sigma(e) = e - \sigma(e) + e\sigma(e) + 1 + \lambda^{-1}u - 2e \\ &= 1 - e - \sigma(e) + e\sigma(e) + \lambda^{-1}u = (1 - e)(1 - \sigma(e)) + \lambda^{-1}u = \lambda^{-1}u. \end{aligned}$$

If $\operatorname{char} F = 2$, the above equalities together with $(1 - e')(1 - \sigma(e')) = 0$ imply that $e'\sigma(e') = 1 + \lambda^{-1}u$. Thus,

$$\dim_F e'\sigma(e')A = \dim_F(1 + \lambda^{-1}u)A = \dim_F(\lambda + u)A = \frac{1}{2}\dim_F A.$$

If $\operatorname{char} F \neq 2$, the element $h := e' - \frac{1}{2}e'\sigma(e')$ is a hyperbolic idempotent with respect to σ. We also have $h - \sigma(h) = e' - \sigma(e') = \lambda^{-1}u$, completing the proof. □

Proposition 5.2. *If $u \in \mathrm{Alt}(A,\sigma)$, then there exists a σ-invariant split quaternion F-subalgebra of A containing u, except for the case where $\operatorname{char} F = 2$, $\lambda = 0$ and σ is orthogonal. In this case there is no σ-invariant quaternion subalgebra $Q \subseteq A$ with $u \in Q$.*

Proof. Suppose first that $\lambda \neq 0$. If $\operatorname{char} F \neq 2$, by (5.1 (3)) there exists a hyperbolic idempotent $h \in A$ such that $\lambda^{-1}u = h - \sigma(h)$ and the result follows from (4.1). If $\operatorname{char} F = 2$, by (5.1 (3)) there exists a metabolic idempotent $e \in A$ such that $\lambda^{-1}u = e - \sigma(e)$ and $\dim_F e\sigma(e)A = \frac{1}{2}\dim_F A$. Hence the conclusion follows from (4.4).

Now, let $\lambda = 0$. The hypotheses imply that either $\operatorname{char} F \neq 2$ or σ is not orthogonal. In either case, by (2.1) there exists $x \in A$ such that $x + \sigma(x) = 1$. Using (5.1 (2)) with $\lambda = 0$, one can find a metabolic idempotent $e \in A$ such that $eu = u$ and $ue = 0$. Then $h := e - ex\sigma(e)$ is a hyperbolic idempotent with respect to σ. As $ue = 0$ we have $uh = 0$. Also, the relations $\sigma(u) = -u$ and

$ue = 0$ imply that $\sigma(e)u = 0$, hence $hu = eu = u$. Hence the result follows from (4.2).

Finally, consider the exceptional case, i.e., suppose that $\operatorname{char} F = 2$, $\lambda = 0$ and σ is orthogonal. If there exists a σ-invariant quaternion subalgebra Q of A containing u, then $u \in \operatorname{Alt}(Q, \sigma|_Q)$ by [9, (3.5)]. By dimension count we obtain $\operatorname{Alt}(Q, \sigma|_Q) = Fu$. But this implies that $\operatorname{Alt}(Q, \sigma|_Q)$ does not contain any invertible element, contradicting [7, (2.8 (2))]. □

Theorem 5.3. *If $\sigma(u) = -u$, then there exists a σ-invariant split quaternion F-subalgebra of A containing u, except for the case where* $\operatorname{char} F = 2$ *and σ is orthogonal.*

Proof. As observed in [7, p. 14] and [7, (2.17)], if $\operatorname{char} F \neq 2$ or σ is unitary, the equality $\sigma(u) = -u$ implies that $u \in \operatorname{Alt}(A, \sigma)$ and the result follows from (5.2). Suppose that $\operatorname{char} F = 2$ and σ is symplectic. Replacing u with $u+1$ if necessary, we may assume that $\lambda \neq 0$ (note that since $\operatorname{char} F = 2$, this replacement does not change the assumptions $\sigma(u) = -u$ and $\dim_F(\lambda + u)A = \frac{1}{2}\dim_F A$). If $u \in \operatorname{Alt}(A, \sigma)$, the conclusion follows from (5.2), hence suppose that $u \notin \operatorname{Alt}(A, \sigma)$. Set $\tau = \operatorname{Int}(u) \circ \sigma$. By [7, (2.7 (3))], τ is an orthogonal involution on A. Since σ is symplectic, by [7, (2.6 (2))] we have $1 \in \operatorname{Alt}(A, \sigma)$. Hence [7, (2.7 (2))] implies that $u \in \operatorname{Alt}(A, \tau)$. By (5.2), there exists a τ-invariant quaternion subalgebra $Q \subseteq A$ containing u. As $u \in Q$ and $\tau(Q) = Q$, we get $\sigma(Q) = u^{-1}\tau(Q)u = Q$, completing the proof. □

Lemma 5.4. *Let* $\operatorname{char} F = 2$ *and let (Q, τ) be a quaternion algebra with orthogonal involution over F. For every symmetric element $x \in Q$ with $x^2 \in F$, there exists $\alpha \in F$ such that $x + \alpha \in \operatorname{Alt}(Q, \tau)$.*

Proof. By [8, (6.1)], Q is generated as an F-algebra by two elements $w, v \in Q$ satisfying

$$w^2 \in F^\times, \quad v^2 \in F, \quad wv + vw = 1, \quad \text{and} \quad \tau(v) + v = w^{-1}.$$

Hence, $\operatorname{Alt}(Q, \tau) = Fw$ and $\operatorname{Sym}(Q, \tau) = F + Fw + Fwv$. Since $x \in \operatorname{Sym}(Q, \tau)$ and $x^2 \in F$, we have $x = \alpha + \beta w$ for some $\alpha, \beta \in F$. Thus, $x + \alpha = \beta w \in \operatorname{Alt}(Q, \tau)$. □

We now consider the exceptional case in (5.3).

Proposition 5.5. *Suppose that* $\operatorname{char} F = 2$*, σ is orthogonal and $\sigma(u) = u$. There exists a σ-invariant quaternion subalgebra of A containing u if and only if $u + \alpha \in \operatorname{Alt}(A, \sigma)$ for some $\alpha \in F$ with $\alpha \neq \lambda$.*

Proof. Suppose that $u \in Q$ for some σ-invariant quaternion subalgebra Q of A. As $\sigma(u) = u$, by (5.4) there exists $\alpha \in F$ such that $u + \alpha \in \operatorname{Alt}(Q, \sigma|_Q) \subseteq \operatorname{Alt}(A, \sigma)$. Since $(u + \alpha)^2 = \lambda^2 + \alpha^2$, using (5.2) we obtain $\alpha \neq \lambda$. To prove the converse, let $u' = u + \alpha \in \operatorname{Alt}(A, \sigma)$ and $\lambda' = \alpha + \lambda$. Then $u'^2 = \lambda'^2 \in F^{\times 2}$ and $\dim_F(u' + \lambda')A = \dim_F(u + \lambda)A = \frac{1}{2}\dim_F A$. The conclusion therefore follows from (5.2). □

We conclude with an example of a symmetric element u in a central simple algebra with involution (A, σ) satisfying $u^2 = \lambda^2 \neq 0$ and $\dim_F(u + \lambda)A = \frac{1}{2}\dim_F A$, which is not contained in any σ-invariant quaternion subalgebra of A.

Example 5.6. Suppose that $\operatorname{char} F = 2$, $\lambda \neq 0$ and $(A, \sigma) = (M_4(F), t)$. Let

$$u = \begin{pmatrix} \lambda & 0 & \lambda & \lambda \\ 0 & \lambda & \lambda & \lambda \\ \lambda & \lambda & 0 & \lambda \\ \lambda & \lambda & \lambda & 0 \end{pmatrix}.$$

Then u is a symmetric matrix, $u^2 = \lambda^2 \in F^2$ and $\dim_F(u + \lambda)A = \frac{1}{2} \dim_F A$. Since $\operatorname{Alt}(A, \sigma)$ consists of all symmetric matrices with zero diagonal, we get $u + \alpha \notin \operatorname{Alt}(A, \sigma)$ for every $\alpha \in F$. Thus, by (5.5) there is no σ-invariant quaternion subalgebra of A containing u.

References

[1] D. Barry, Power-central elements in tensor products of symbol algebras. LAG preprint server, http://www.math.uni-bielefeld.de/LAG/man /516.html.

[2] E. Bayer-Fluckiger, D. B. Shapiro, J.-P. Tignol, Hyperbolic involutions. *Math. Z.* **214** (1993), no. 3, 461–476.

[3] G. Berhuy, C. Frings, J.-P. Tignol, Galois cohomology of the classical groups over imperfect fields. *J. Pure Appl. Algebra* **211** (2007), no. 2, 307–341.

[4] Đoković, Dragomir Ž. Inner derivations of division rings and canonical Jordan form of triangular operators. *Proc. Amer. Math. Soc.* **94** (1985), no. 3, 383–386.

[5] A. Dolphin, Metabolic involutions. *J. Algebra* **336** (2011), 286–300.

[6] P. K. Draxl, Skew fields. London Mathematical Society Lecture Note Series, 81. *Cambridge University Press, Cambridge*, 1983.

[7] M.-A. Knus, A. S. Merkurjev, M. Rost, J.-P. Tignol, *The book of involutions.* American Mathematical Society Colloquium Publications, 44. American Mathematical Society, Providence, RI, 1998.

[8] M. G. Mahmoudi, A.-H. Nokhodkar, Involutions of a Clifford algebra induced by involutions of orthogonal group in characteristic 2. *Comm. Algebra* **43** (2015), no. 9, 3898–3919.

[9] M. G. Mahmoudi, A.-H. Nokhodkar, On split products of quaternion algebras with involution in characteristic two. *J. Pure Appl. Algebra* **218** (2014), no. 4, 731–734.

A.-H. Nokhodkar, `a.nokhodkar@kashanu.ac.ir`
Department of Pure Mathematics, Faculty of Science, University of Kashan, P. O. Box 87317-53153, Kashan, Iran.